

SUPA TRACE[®] ADVANCE

NPKS 3-0-0-4

+ Iron, Zinc, Copper, Manganese, Boron and Molybdenum



Supa Trace[®] Advance is complexed for maximum bioavailability, with all essential trace elements required for your crop to reach its full potential

BENEFITS OF SUPA TRACE[®] ADVANCE

- Replenishes the plant with all required trace elements that are vital for healthy growth and yield maximization.
- Complexation of the nutrients increases plant availability and rate of uptake.
- Contains nitrogen which facilitates trace element uptake by plants.
- A well balanced mix of essential trace elements which maintain plant health and reduce the potential for deficiencies.
- Improves overall plant development and utilisation of major nutrients applied.
- Eliminates trace element deficiency syndrome.

THE IMPORTANCE OF TRACE ELEMENTS

Many trace elements function as essential part of enzymes in the cell. Important enzymes consist of proteins which attach to co-enzymes. The control of cellular processes through chemical reactions is performed through enzymes.

Zinc forms many enzymes, which maintains respiration, protein synthesis, photosynthesis and production of auxins.

Copper is crucial to several enzyme systems. It is involved in cell wall formation, electron transport and oxidation reactions. Copper also affects the formation and chemical composition of cell walls.

Manganese is an enzyme activator which helps with nitrate assimilation. It is primarily involved with photosynthesis and chlorophyll production.

Iron is required to produce chlorophyll and to activate several enzymes, especially those involved in the oxidation / reduction processes of photosynthesis and respiration.

Boron is needed for sugar movement within the plant, as well as formation of new cells at growing points. Boron also affects pollination and seed development.

Molybdenum is essential for the chemical changes involved with nitrogen assimilation during the conversion of nitrate nitrogen to ammonium inside the plant. It is important for chlorophyll and enzyme formation.

SUPA TRACE[®] ADVANCE

CHARACTERISTICS: pH: 1.0 – 2.0; Specific Gravity: 1.27 – 1.30

AUS Analysis W/V%: 2.5% N, 4.4% S, 1.3% Fe, 1.5% Zn, 0.5% Cu, 1.1% Mn, 1.1% Mg, 0.5% B, 0.01% Mo

International Analysis W/W%: 2% N, 3.5% S, 1% Fe, 1.2% Zn, 0.4% Cu, 0.9% Mn, 0.9% Mg, 0.4% B, 0.01% Mo

APPLICATION

BROADACRE: Such as Barley, Canola, Cotton, Grain legumes, Maize, Oats, Rice, Sorghum, Triticale, Wheat & Pasture crops. **Foliar: 2 – 4 L/ha** in a minimum of 50 - 80 L final spray volume. Best applied at 3 – 4 true leaf, may be used at other growth stages. Use low rate for low density crops (140 – 160 plants m²) & higher rate for high yielding crops. Apply in 50 – 80 L of water / ha. **Use higher dilutions in temperatures > 28°C. Apply prior to flowering in Canola. Aerial application: use maximum practicable water rates.**

CUT FLOWERS & ORNAMENTALS OPEN FIELD: Such as Carnations, Gypsophilla, Roses & Statice. **Fertigation: 5 – 7 L/ha.** Apply regularly to replenish nutrients and maintain colour.

DECIDUOUS TREE CROPS: Such as Apple, Almond, Cherry, Nectarine, Peach, Pear, Pistachio and Walnut. **Foliar: 3 – 5 L/ha** in a minimum of 200 - 300L final spray volume. **Fertigation: 5 – 7 L/ha.** Apply to newly hardened spring flush or during active growing period and post-harvest. **DO NOT apply to Stone fruit in season as phytotoxicity will occur. Dormancy spray only.**

EVERGREEN TREE CROPS: Such as Avocado, Citrus, Macadamia, Mangoes, Lychee. **Foliar: 3 – 5 L/ha** in a minimum of 400 - 800L final spray volume. **Fertigation: 5 – 7 L/ha.** Apply to newly hardened spring flush or during active growing period and post-harvest.

FRUITING VEGETABLES: Such as Capsicum, Cucurbits, Eggplant, Tomatoes, Watermelons, Pumpkins. **Foliar: 3 – 5 L/ha** in a minimum of 600 - 1000L final spray volume. Apply as required to maintain trace element levels. Typically, at 14 – 21 day intervals. **Fertigation: 5 – 7 L/ha.** Apply regularly to replenish nutrients.

LEAFY VEGETABLES: Such as Endive, Fennel Lettuce, Broccoli, Cabbage, Cauliflower, Kale and Herbs. **Foliar: 2 – 4 L/ha** in a minimum of 400 - 800L final spray volume. Apply as required to maintain trace element levels. **Fertigation: 5 – 7 L/ha.** Apply regularly to replenish nutrients.

ROOT VEGETABLES: Such as Beetroot, Carrot, Leek, Onion, Potato, Radish, Sweet Potato. **Foliar: 2 – 4 L/ha** in a minimum of 400 - 800L final spray volume. **Fertigation: 5 – 7 L/ha.** Apply as required to maintain trace element levels. Apply with compatible crop protection sprays.

VINE and BERRY CROPS: Such as Blueberry, Strawberry, Raspberry, Wine and Table Grapes. **Foliar: 2 – 4 L/ha** in a minimum of 500 - 1000L final spray volume. **Fertigation: 5 – 7 L/ha.** Maximum of 2 applications may be required post-flowering up to veraison.

Fertigation rates are dependent on seasonal nutrient demand. Aerial applications: use maximum practical water rates.

Note when applying in alkaline conditions (water or soil) ensure product to water ratio is 1:100. Agitate contents well prior to application.

DO NOT apply to fruit with copper residues.

The information contained in this Product Information Sheet in respect of the "Product" is indicative only and should not be relied upon as advice or a recommendation. While this Information Sheet has been prepared in good faith, Agrichem does not warrant the accuracy of this information. You use the information at your own risk and should rely on your own independent inquiries and assessments. With the exception of the consumer guarantees provided by the Australian Consumer Law (ACL), all conditions and warranties implied in respect of any information or advice provided by Agrichem about the Product are excluded, and Agrichem does not accept any liability whatsoever (including through misrepresentation or negligence), incurred in connection with your use or reliance upon this Information Sheet. If liability under the ACL cannot be excluded but the Product the subject of the Information Sheet is NOT used for personal, domestic or household use or consumption, Agrichem may (at its election) limit its liability to replacement of the Product, or payment of the cost of acquiring the Product. You must not reproduce this information sheet without written consent from Agrichem®.

NOTE: The suggested rates of application of the Product are designed for typical Australian conditions and should be used as a guide only. Each farmer's climatic conditions, water quality, soil types, application processes and practices may differ and therefore necessitate corrections to ensure optimum results. Good agricultural practice requires that application be avoided under extreme weather conditions such as temperatures over 28°C, high humidity, frost, rain etc. It is recommended that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total spray. Where possible, it is recommended that regular leaf tests are conducted to determine actual plant nutrient availability during each growth cycle. Soil tests at least once per year are essential.