

Biological Soil conditioner.



A unique bio soil conditioner that reduces salinity, helps improve soil structure and aeration. Plants exhibit better nutrient uptake and reduction of stress.

## **BENEFITS OF AGRI TIGER<sup>®</sup>**

- Helps release micro and macro elements from its insoluble state form in irrigation water and soil.
- Helps plants overcome high salinity conditions in soils.
- Reduces the impact of high soil EC on plant growth.
- Improves nutrient uptake.
- Improves nutrient efficciency of applied nutrients in all soil types.
- Improves root development.

Agri Tiger can be applied on;

- I. Fatigued soils
- 2. Soils with high sodium levels
- 3. Residue (chemical) contaminated soils
- 4. Compacted soils
- 5. Soils with a long cropping history & related fertiliser applications

## **IMPORTANCE OF AGRI TIGER®**

Agri Tiger is a unique highly active biological soil conditioner. It is made from a primordial cluster of microbes, which preceded the development of higher plants and animals.

Agri Tiger microbes are indifferent to the percent (%) of organic matter in the soil and have no symbiotic relation with plants. It acts independently.

Agri Tiger is critical in re-establishment of soil biology in over exploited farming fields. Locked nutrient elements become more available to plants and after a period of 2- month one can observe significant changes in the plants performance. You will also note that in just a few weeks soils structure is enhanced with noticeable improvement in soil friability & reduction in soil compaction.



## AGRI TIGER<sup>®</sup> CHARACTERISTICS: pH: 2.0 - 5.0; Specific Gravity: 1.04 - 1.1

**AUS Analysis W/V%:** 0.62% Mg, 0.052% Zn. **International Analysis W/W%:** 0.6% Mg, 0.05 % Zn.

## **APPLICATION**

Agri Tiger is designed for use on all crops, ensuring effective and beneficial results for every cultivation.

Recommended rate is 20 Lit / Ha drench once a year and can be applied at any time (field preparation, during the crop growth). If the water source has high levels of sodium we recommend 2 applications a year.

The information contained in this Product Information Sheet in respect of the "Product" is indicative only and should not be relied upon as advice or a recommendation. While this Information Sheet has been prepared in good faith, Agrichem does not warrant the accuracy of this information.You use the information at your own risk and should rely on your own

While this Information Sheet has been prepared in good faith, Agrichem does not warrant the accuracy of this information. You use the information at your own risk and should rely on your own independent inquiries and assessments. With the exception of the consumer guarantees provided by the Australian Consumer Law (ACL), all conditions and warranties implied in respect of any information or advice provided by Agrichem about the Product are excluded, and Agrichem does not accept any liability whatsoever (including through misrepresentation or negligence), incurred in connection with your use or reliance upon this Information Sheet. If liability under the ACL cannot be excluded but the Product the subject of the Information Sheet is NOT used for personal, domestic or household use or consumption, Agrichem may (at its election) limit its liability to replacement of the Product, or payment of the cost of acquiring the Product. You must not reproduce this information sheet without written consent from Agrichem@.

NOTE: The suggested rates of application of the Product are designed for typical Australian conditions and should be used as a guide only. Each farmer's climatic conditions, water quality, soil types, application processes and practices may differ and therefore necessitate corrections to ensure optimum results. Good agricultural practice requires that application be avoided under extreme weather conditions such as temperatures over 28<sup>°C</sup>, high humidity, frost, rain etc. It is recommended that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total spray. Where possible, it is recommended that regular leaf tests are conducted to determine actual plant nutrient availability during each growth cycle. Soil tests at least once per year are essential.